

A TeV-scale model for neutrino mass, DM and baryon asymmetry

Mayumi Aoki

Department of Physics, Tohoku University, Aramaki, Aoba, Sendai, Miyagi 980-8578, JAPAN

E-mail: mayumi@tuhep.phys.tohoku-u.ac.jp

Shinya Kanemura*

Department of Physics, Univeristy of Toyama, 3190 Gofuku, Toyama 930-8555, JAPAN

E-mail: kanemu@sci.u-toyama.ac.jp

Osamu Seto

Instituto de Física Teórica UAM/CSIC, Universidad Autónoma de Madrid, Cantoblanco, Madrid 28049, Spain

E-mail: osamu.seto@uam.es

We discuss a model which would explain neutrino oscillation, dark matter, and baryon asymmetry of the Universe simultaneously by the physics at TeV scales. Tiny neutrino masses are generated at the three loop level due to the exact Z_2 symmetry, by which the stability of the dark matter candidate is also guaranteed. The extra Higgs doublet is required not only for the tiny neutrino masses but also for successful electroweak baryogenesis. The model provides various discriminative predictions especially in charged Higgs phenomenology.

Prospects for Charged Higgs Discovery at Colliders

September 16-19 2008

Uppsala, Sweden

*Speaker.

1. Introduction

Today, we have clear motivations to consider a model beyond the standard model (SM). The data indicate that neutrinos have tiny masses and mix with each other[1]. Cosmological observations have revealed that the energy density of dark matter (DM) in the Universe dominates that of baryonic matter[2]. Asymmetry of matter and anti-matter in our Universe has been addressed for several decades[3].

In this talk, we discuss a model which would explain neutrino oscillation, origin of DM and baryon asymmetry simultaneously by an extended Higgs sector with TeV-scale right-handed (RH) neutrinos[4]. Tiny neutrino masses are generated at the three loop level due to an exact discrete symmetry, by which tree-level Yukawa couplings of neutrinos are prohibited. The lightest neutral odd state under the discrete symmetry is a candidate of DM. Baryon asymmetry can be generated at the electroweak phase transition (EWPT) by additional CP violating phases in the Higgs sector[5]. In this framework, a successful model can be built without contradiction of the current data.

Original idea of generating tiny neutrino masses via the radiative effect has been proposed by Zee[6]. The extension with a TeV-scale RH neutrino has been discussed in Ref. [7], where neutrino masses are generated at the three-loop level due to the exact Z_2 parity, and the Z_2 -odd RH neutrino is a candidate of DM. This has been extended with two RH neutrinos to describe the neutrino data[8]. Several models with adding baryogenesis have been considered in Ref. [9]. The following advantages would be in the present model: (a) all mass scales are at most at the TeV scale without large hierarchy, (b) physics for generating neutrino masses is connected with that for DM and baryogenesis, (c) the model parameters are strongly constrained by the current data, so that the model provides discriminative predictions which can be tested at future experiments.


2. Model

We introduce two scalar isospin doublets with hypercharge 1/2 (Φ_1 and Φ_2), charged singlets ($S^\pm$), a real scalar singlet (η) and two generation isospin-singlet RH neutrinos (N_R^α with $\alpha = 1, 2$). As in Ref. [7] we impose an exact Z_2 symmetry to generate tiny neutrino masses at the three-loop level, which we refer as Z_2 . We assign Z_2 -odd charge to $S^\pm$, η and N_R^α , while ordinary gauge fields, quarks and leptons and Higgs doublets are Z_2 even. Introduction of two Higgs doublets would cause a dangerous flavor changing neutral current. To avoid this in a natural way, we impose another discrete symmetry ($\tilde{Z}_2$) that is softly broken[10]. From a phenomenological reason discussed later, we assign $\tilde{Z}_2$ charges such that only Φ_1 couples to leptons whereas Φ_2 does to quarks;

$$\mathcal{L}_Y = -y_{e_l} \bar{L}^i \Phi_1 e_R^i - y_{u_l} \bar{Q}^i \tilde{\Phi}_2 u_R^i - y_{d_l} \bar{Q}^i \Phi_2 d_R^i + \text{h.c.}, \quad (2.1)$$

where Q^i (L^i) is the ordinary i -th generation left-handed (LH) quark (lepton) doublet, and u_R^i and d_R^i (e_R^i) are RH-singlet up- and down-type quarks (charged leptons), respectively. We summarize the particle properties under Z_2 and $\tilde{Z}_2$ in Table 1. Notice that the Yukawa coupling in Eq. (2.1) is different from that in the minimal supersymmetric SM (MSSM)[11]. In addition to the usual potential of the two Higgs doublet model (THDM) with the $\tilde{Z}_2$ parity and that of the Z_2 -odd scalars,

	Q^i	u_R^i	d_R^i	L^i	e_R^i	Φ_1	Φ_2	$S^\pm$	η	N_R^α
Z_2 (exact)	+	+	+	+	+	+	+	-	-	-
$\tilde{Z}_2$ (softly broken)	+	-	-	+	+	+	-	+	-	+

Table 1: Particle properties under the discrete symmetries.**Figure 1:** The diagrams for generating tiny neutrino masses.

we have the interaction terms between Z_2 -even and -odd scalars:

$$\mathcal{L}_{int} = - \sum_{a=1}^2 \left(\rho_a |\Phi_a|^2 |S|^2 + \sigma_a |\Phi_a|^2 \frac{\eta^2}{2} \right) - \sum_{a,b=1}^2 \{ \kappa \varepsilon_{ab} (\Phi_a^c)^\dagger \Phi_b S^- \eta + \text{h.c.} \}, \quad (2.2)$$

where ε_{ab} is the anti-symmetric tensor with $\varepsilon_{12} = 1$. The mass term and the interaction for N_R^α are given by

$$\mathcal{L}_{Y_N} = \sum_{\alpha=1}^2 \left\{ \frac{1}{2} m_{N_R^\alpha} \overline{N_R^{\alpha c}} N_R^\alpha - h_i^\alpha \overline{(e_R^i)^c} N_R^\alpha S^- + \text{h.c.} \right\}. \quad (2.3)$$

Although the CP violating phase in the Lagrangian is crucial for successful baryogenesis at the EWPT[5], it does not much affect the following discussions. Thus, we neglect it for simplicity. We later give a comment on the case with the non-zero CP-violating phase.

As Z_2 is exact, the even and odd fields cannot mix. Mass matrices for the Z_2 even scalars are diagonalized as in the usual THDM by the mixing angles α and β , where α diagonalizes the CP-even states, and $\tan \beta = \langle \Phi_2^0 \rangle / \langle \Phi_1^0 \rangle$ [11]. The Z_2 even physical states are two CP-even (h and H), a CP-odd (A) and charged ($H^\pm$) states. We here define h and H such that h is always the SM-like Higgs boson when $\sin(\beta - \alpha) = 1$.

3. Neutrino Mass, Dark Matter, and Strongly 1st-Order Phase Transition

The LH neutrino mass matrix M_{ij} is generated by the three-loop diagrams in Fig. 1. The absence of lower order loop contributions is guaranteed by Z_2 . $H^\pm$ and e_R^i play a crucial role to connect LH neutrinos with the one-loop sub-diagram by the Z_2 -odd states. We obtain

$$M_{ij} = \sum_{\alpha=1}^2 C_{ij}^\alpha F(m_{H^\pm}, m_{S^\pm}, m_{N_R^\alpha}, m_\eta), \quad (3.1)$$

Set	h_e^1	h_e^2	h_μ^1	h_μ^2	h_τ^1	h_τ^2	$B(\mu \rightarrow e\gamma)$
A	2.0	2.0	-0.019	0.042	-0.0025	0.0012	6.9×10^{-12}
B	2.2	2.2	0.0085	0.038	-0.0012	0.0021	6.1×10^{-12}

Table 2: Values of h_i^α for $m_{H^\pm}(m_{S^\pm}) = 100(400)\text{ GeV}$, $m_\eta = 50\text{ GeV}$, $m_{N_R^1} = m_{N_R^2} = 3.0\text{ TeV}$ for the normal hierarchy. For Set A (B), $\kappa \tan \beta = 28(32)$ and $U_{e3} = 0(0.18)$. Predictions on the branching ratio of $\mu \rightarrow e\gamma$ are also shown.




Figure 2: [Left figure] The relic abundance of η . [Right figure] The region of strong first order EWPT. Deviations from the SM value in the hhh coupling are also shown.

where $C_{ij}^\alpha = 4\kappa^2 \tan^2 \beta (y_{e_i}^{\text{SM}} h_i^\alpha) (y_{e_j}^{\text{SM}} h_j^\alpha)$ with $y_{e_i}^{\text{SM}} = \sqrt{2} m_{e_i} / v$ and $v \simeq 246\text{ GeV}$. The factor of the three-loop integral function $F(m_{H^\pm}, m_{S^\pm}, m_{N_R}, m_\eta)$ includes the suppression factor of $1/(16\pi^2)^3$, whose typical size is $\mathcal{O}(10^4)\text{ eV}$. Magnitudes of $\kappa \tan \beta$ as well as F determine the universal scale of M_{ij} , whereas variation of h_i^α ($i = e, \mu, \tau$) reproduces the mixing pattern indicated by the neutrino data[1].

Under the *natural* requirement $h_e^\alpha \sim \mathcal{O}(1)$, and taking the $\mu \rightarrow e\gamma$ search results into account[12], we find that $m_{N_R^\alpha} \sim \mathcal{O}(1)\text{ TeV}$, $m_{H^\pm} \lesssim \mathcal{O}(100)\text{ GeV}$, $\kappa \tan \beta \gtrsim \mathcal{O}(10)$, and $m_{S^\pm}$ being several times 100 GeV. On the other hand, the LEP direct search results indicate $m_{H^\pm}$ (and $m_{S^\pm}$) $\gtrsim 100\text{ GeV}$ [1]. In addition, with the LEP precision measurement for the ρ parameter, possible values uniquely turn out to be $m_{H^\pm} \simeq m_H$ (or m_A) $\simeq 100\text{ GeV}$ for $\sin(\beta - \alpha) \simeq 1$. Thanks to the Yukawa coupling in Eq. (2.1), such a light $H^\pm$ is not excluded by the $b \rightarrow s\gamma$ data[13]. Since we cannot avoid to include the hierarchy among y_i^{SM} , we only require $h_i^\alpha y_i \sim \mathcal{O}(y_e) \sim 10^{-5}$ for values of h_i^α . Our model turns out to prefer the normal hierarchy scenario. Several sets for h_i^α are shown in Table 2 with the predictions on the branching ratio of $\mu \rightarrow e\gamma$ assuming the normal hierarchy.

The lightest Z_2 -odd particle is stable and can be a candidate of DM if it is neutral. In our model, N_R^α must be heavy, so that the DM candidate is identified as η . When η is lighter than the W boson, η dominantly annihilates into $b\bar{b}$ and $\tau^+\tau^-$ via tree-level s -channel Higgs (h and H) exchange diagrams, and into $\gamma\gamma$ via one-loop diagrams. From their summed thermal averaged annihilation rate $\langle \sigma v \rangle$, the relic mass density $\Omega_\eta h^2$ is evaluated. Fig. 2(Left) shows $\Omega_\eta h^2$ as a function of m_η . Strong annihilation can be seen near $50\text{ GeV} \simeq m_H/2$ ($60\text{ GeV} \simeq m_h/2$) due to the resonance of H (h) mediation. The data ($\Omega_{\text{DM}} h^2 \simeq 0.11$ [2]) indicate that m_η is around 40-65 GeV.

The model satisfies the necessary conditions for baryogenesis[3]. Especially, departure from thermal equilibrium can be realized by the strong first order EWPT. The free energy is given at a high temperature T by

$$V_{eff}[\phi, T] = D(T^2 - T_0^2)\phi^2 - ET\phi^3 + \frac{\lambda_T}{4}\phi^4 + \dots, \quad (3.2)$$

where ϕ is the order parameter. A large value of the coefficient E is crucial for the strong first order EWPT with keeping $m_h \lesssim 120$ GeV. For sufficient sphaleron decoupling in the broken phase, it is required that[14]

$$\frac{\phi_c}{T_c} \left(\simeq \frac{2E}{\lambda_{T_c}} \right) \gtrsim 1, \quad (3.3)$$

where $\phi_c (\neq 0)$ and T_c are the critical values of ϕ and T at the EWPT. In Fig. 2(Right), the allowed region under the condition of Eq. (3.3) is shown. The condition is satisfied when $m_{S^\pm} \gtrsim 350$ GeV for $m_A \gtrsim 100$ GeV, $m_h \simeq 120$ GeV, $m_H \simeq m_{H^\pm} (\simeq M) \simeq 100$ GeV and $\sin(\beta - \alpha) \simeq 1$.

4. Phenomenology

A successful scenario which can simultaneously solve the above three issues under the data[1, 12, 13] would be

$$\begin{aligned} \sin(\beta - \alpha) &\simeq 1, & (\kappa \tan \beta) &\simeq 30, \\ m_h &= 120 \text{ GeV}, & m_H &\simeq m_{H^\pm} (\simeq M) \simeq 100 \text{ GeV}, \\ m_A &\gtrsim 100 \text{ GeV}, & m_{S^\pm} &\sim 400 \text{ GeV}, \\ m_\eta &\simeq 40 - 65 \text{ GeV}, & m_{N_R^1} &\simeq m_{N_R^2} \simeq 3 \text{ TeV}. \end{aligned} \quad (4.1)$$

This is realized without assuming unnatural hierarchy among the couplings. All the masses are between $\mathcal{O}(100)$ GeV and $\mathcal{O}(1)$ TeV. As they are required by the data, the model has a predictive power. We note that the masses of A and H can be exchanged with each other.

We outline phenomenological predictions in the scenario in (4.1) in order. The detailed analysis is shown elsewhere[15]. (I) h is the SM-like Higgs boson, but decays into $\eta\eta$ when $m_\eta < m_h/2$. The branching ratio is about 36% (25%) for $m_\eta \simeq 45$ (55) GeV. This is related to the DM abundance, so that our DM scenario is testable at the LHC. (II) η is potentially detectable by direct DM searches[16], because η can scatter with nuclei via the scalar exchange[17]. (III) For successful baryogenesis, the hhh coupling has to deviate from the SM value by more than 10-20 % [18] (see Fig. 2), which can be tested at the International Linear Collider (ILC)[19]. (IV) H (or A) can predominantly decay into $\tau^+\tau^-$ instead of $b\bar{b}$ for $\tan \beta \gtrsim 3$. When A (or H) is relatively heavy it can decay into $H^\pm W^\mp$ and HZ (or AZ). (V) the scenario with light $H^\pm$ and H (or A) can be directly tested at the LHC via $pp \rightarrow W^* \rightarrow HH^\pm$ and $AH^\pm$ [20]. (VI) $S^\pm$ can be produced in pair at the LHC (the ILC)[21], and decay into $\tau^\pm \nu \eta$. The signal would be a hard hadron pair[22] with a large missing energy. (VII) The couplings h_i^α cause lepton flavor violation such as $\mu \rightarrow e\gamma$ which would provide information on $m_{N_R^\alpha}$ at future experiments.

Finally, we comment on the case with the CP violating phases. Our model includes the THDM, so that the same discussion can be applied in evaluation of baryon number at the EWPT[5]. The mass spectrum would be changed to some extent, but most of the features discussed above should be conserved with a little modification.

5. Summary

We have discussed the model with the extended Higgs sector and TeV-scale RH neutrinos, which would explain neutrino mass and mixing, DM and baryon asymmetry by the TeV scale physics without less fine tuning. It gives specific predictions on many low energy phenomena. In particular, charged Higgs phenomenology in this model is completely different from that in the MSSM, so that it is testable at the LHC and the ILC.

References

- [1] W. M. Yao, et al., [Particle Data Group] J. Phys. G **33** (2006) 1.
- [2] E. Komatsu, et al., (WMAP Collaboration), arXiv:0803.0547 [astro-ph].
- [3] A. D. Sakharov, Pisma Zh. Eksp. Teor. Fiz. **5**, 32 (1967).
- [4] M. Aoki, S. Kanemura and O. Seto, arXiv:0807.0361 [hep-ph].
- [5] J. M. Cline, K. Kainulainen and A. P. Vischer, Phys. Rev. D **54**, 2451 (1996); L. Fromme, S. J. Huber and M. Seniuch, JHEP **0611**, 038 (2006).
- [6] A. Zee, Phys. Lett. B **93**, 389 (1980) [Erratum-ibid. B **95**, 461 (1980)]; A. Zee, Phys. Lett. B **161**, 141 (1985).
- [7] L. M. Krauss, S. Nasri and M. Trodden, Phys. Rev. D **67**, 085002 (2003).
- [8] K. Cheung and O. Seto, Phys. Rev. D **69**, 113009 (2004).
- [9] E. Ma, Phys. Rev. D **73**, 077301 (2006); J. Kubo, E. Ma and D. Suematsu, Phys. Lett. B **642**, 18 (2006); T. Hambye, K. Kannike, E. Ma and M. Raidal, Phys. Rev. D **75**, 095003 (2007); K. S. Babu and E. Ma, arXiv:0708.3790 [hep-ph]; N. Sahu and U. Sarkar, arXiv:0804.2072 [hep-ph].
- [10] S. L. Glashow and S. Weinberg, Phys. Rev. D **15**, 1958 (1977); V. D. Barger, J. L. Hewett and R. J. N. Phillips, Phys. Rev. D **41**, 3421 (1990).
- [11] J. F. Gunion, et al., “*The Higgs Hunters’s Guide*” (Addison Wesley, 1990).
- [12] A. Baldini, Nucl. Phys. Proc. Suppl. **168**, 334 (2007).
- [13] E. Barberio, et al., [Heavy Flavor Averaging Group (HFAG)], arXiv:hep-ex/0603003.
- [14] G. D. Moore, Phys. Lett. B **439**, 357 (1998); Phys. Rev. D **59**, 014503 (1998).
- [15] M. Aoki, S. Kanemura and O. Seto, in preparation.
- [16] Y. D. Kim, Phys. Atom. Nucl. **69**, 1970 (2006); D. S. Akerib, et al., [CDMS Collaboration], Phys. Rev. Lett. **96**, 011302 (2006).
- [17] J. McDonald, Phys. Rev. D **50**, 3637 (1994); for a recent study, see *e.g.*, H. Sung Cheon, S. K. Kang and C. S. Kim, J. Cosmol. Astropart. Phys. **05** (2008) 004.
- [18] S. Kanemura, Y. Okada and E. Senaha, Phys. Lett. B **606**, 361 (2005).
- [19] M. Battaglia, E. Boos and W. M. Yao, arXiv:hep-ph/0111276; Y. Yasui, et al., arXiv:hep-ph/0211047.
- [20] S. Kanemura and C. P. Yuan, Phys. Lett. B **530**, 188 (2002); Q. H. Cao, S. Kanemura and C. P. Yuan, Phys. Rev. D **69**, 075008 (2004).
- [21] S. Kanemura, T. Kasai, G. L. Lin, Y. Okada, J. J. Tseng and C. P. Yuan, Phys. Rev. D **64**, 053007 (2001).
- [22] B. K. Bullock, K. Hagiwara and A. D. Martin, Phys. Rev. Lett. **67**, 3055 (1991).